

### AMENDMENTS TO THE CLAIMS

There have been no amendments to the claims (except withdrawal of claims 17-24, 26-33, and 38). However, they are presented below for purposes of convenience.

1. (Original) A method of estimating a parameter of a local maxima or minima of a function comprising:
  - performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;
  - deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and
  - deriving an estimate of the parameter from the interpolation offset.
2. (Original) The method of claim 1 wherein the function is a correlation function.
3. (Original) The method of claim 2 wherein the correlation function is derived from a received signal.
4. (Original) The method of claim 2 wherein the second deriving step comprises deriving a parameter bias from the interpolation offset using a pre-existing relationship that is present between these two variables and then deriving an estimate of the parameter from the parameter bias.
5. (Original) The method of claim 4 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a code phase dimension.
6. (Original) The method of claim 5 wherein the parameter bias is a code phase bias.
7. (Original) The method of claim 6 wherein the parameter which is

estimated is location of a peak along the code phase dimension, and an estimate of this parameter is derived from the code phase bias.

8. (Original) The method of claim 4 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a Doppler frequency dimension.

9. (Original) The method of claim 8 wherein the parameter bias is a Doppler frequency bias.

10. (Original) The method of claim 9 wherein the parameter which is estimated is location of a peak of the function along the Doppler frequency dimension, and an estimate of this parameter is derived from the Doppler frequency bias.

11. (Original) The method of claim 5 wherein the parameter bias is a peak energy bias.

12. (Original) The method of claim 11 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak energy bias.

13. Canceled.

14. Canceled.

15. (Original) The method of claim 4 wherein the pre-existing relationship between the interpolation offset and the parameter bias is embodied as a lookup table.

16. (Original) The method of claim 15 wherein the second deriving step comprises directly deriving an estimate of the parameter from the interpolation offset through an access to the lookup table.

17. Withdrawn.
18. Withdrawn.
19. Withdrawn.
20. Withdrawn.
21. Withdrawn.
22. Withdrawn.
23. Withdrawn.
24. Withdrawn.
25. Canceled.
26. Withdrawn.
27. Withdrawn.
28. Withdrawn.
29. Withdrawn.
30. Withdrawn.
31. Withdrawn.

32. Withdrawn.

33. Withdrawn.

34. Canceled.

35. Canceled.

36. Canceled.

37. Canceled.

38. Withdrawn.

39. (Original) A memory tangibly embodying a sequence of software instructions for performing a method of estimating a parameter of a local maxima or minima of a function comprising:  
performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;  
deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and  
deriving an estimate of the parameter from the interpolation offset.

40. (Original) The memory of claim 39 wherein the function is a correlation function.

41. (Original) The memory of claim 40 wherein the correlation function is derived from a received signal.

42. (Original) The memory of claim 40 wherein the second deriving step

comprises deriving a parameter bias from the interpolation offset using a pre-existing relationship which is present between these two variables and deriving an estimate of the parameter from the parameter bias.

43. (Original) The memory of claim 42 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a code phase dimension.

44. (Original) The memory of claim 43 wherein the parameter bias is a code phase bias.

45. (Original) The memory of claim 44 wherein the parameter is location of a peak along the code phase dimension, and an estimate of this parameter is derived from the code phase bias.

46. (Original) The memory of claim 42 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a Doppler frequency dimension.

47. (Original) The memory of claim 46 wherein the parameter bias is a Doppler frequency bias.

48. (Original) The memory of claim 47 wherein the parameter which is estimated is location of a peak of the function along the Doppler frequency dimension, and an estimate of this parameter is derived from the Doppler frequency bias.

49. (Original) The memory of claim 46 wherein the parameter bias is a peak energy bias.

50. (Original) The memory of claim 49 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak

energy bias.

51. (Original) The memory of claim 46 wherein the parameter bias is a peak energy bias.

52. (Original) The memory of claim 51 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak energy bias.

53. (Original) The memory of claim 42 wherein the pre-existing relationship between the interpolation offset and the parameter bias is embodied as a lookup table.

54. (Original) A system comprising a processor and the memory of claim 39, wherein the processor is configured to access and execute the sequence of software instructions tangibly embodied by the memory.

55. (Original) A method of estimating a parameter of a local maxima or minima of a function comprising:

- a step for performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;

- a step for deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and

- a step for deriving an estimate of the parameter from the interpolation offset.